
RSNA Press Release

Nuclear Stress Test Helps Identify Heart Attack Risk in People with Diabetes

Released: June 27, 2019

OAK BROOK, Ill. — Abnormal results on a nuclear stress test are associated with a significantly increased risk of cardiac-related deaths, especially among people with diabetes, according to a multi-center study published in the journal *Radiology: Cardiothoracic Imaging*.

Researchers said the study results support a role for Positron Emission Tomography (PET) Myocardial Perfusion Imaging (MPI) in improving cardiac risk stratification in people with diabetes.



Hicham Skali, M.D., M.Sc.

Diabetes is a devastating worldwide pandemic and a major risk factor for coronary artery disease and heart attack. Accurate cardiac risk assessment of these patients is vital, as it helps guide appropriate and potentially lifesaving treatments.

PET MPI has emerged in recent years as a promising diagnostic and risk assessment tool. The test uses radioisotopes and a special camera to show how well blood is flowing in the heart when it is under stress. Prior research backs the test's use in evaluating individuals with suspected or known coronary heart disease. However, less is known about its value among patients with diabetes.

At A Glance

- A nuclear stress test called PET MPI can help identify diabetic patients who are at greater risk for coronary artery disease and heart attack.
- PET MPI uses radioisotopes and a special camera to show how well blood is flowing in the heart when it is under stress.
- The results showed that an abnormal PET MPI was associated with increased risk of cardiac death in all important clinical subgroups of diabetic patients.

In the largest study of its kind to date, investigators at four centers collected clinical data on stress tests for both diabetic and non-diabetic patients and then followed them to track the occurrence of adverse events like heart attacks. The study group consisted of 7,061 participants, including 1,966 people with diabetes.

The data showed that among diabetic patients an abnormal PET MPI was associated with increased risk of cardiac death in all important clinical subgroups based on age, gender, obesity, or those with prior revascularization procedures like angioplasty. Using the data, the researchers were able to more accurately assess the cardiac risk for a significant proportion of diabetic patients.

"The data from the stress test among diabetic patients actually allowed us to better risk-stratify people in greater than 39 percent of the cases," said study lead author Hicham Skali, M.D., M.Sc., from Brigham and Women's Hospital and Harvard Medical School in Boston. "Patients with diabetes remain at a significantly higher risk of cardiac death compared to patients without diabetes, and the data from a stress test helps us further stratify those at greatest risk."

The results suggest PET MPI could help select which vulnerable patients may need immediate treatment while sparing others from unnecessary procedures.

Among people without diabetes, the study findings echoed those of previous studies that showed men face a significantly higher risk of cardiac death than women. But that wasn't the case among diabetic patients.

"In patients without diabetes, being a woman confers a certain advantage in that their risk of death is much lower, regardless of their stress findings," Dr. Skali said. "However, when you look at patients with diabetes, men and women have relatively the same risk of cardiovascular death, and that risk increases with worsening findings on the PET stress test."

The data also revealed that, even when they had normal PET MPI results, people with diabetes had a similar rate of cardiac death to people without diabetes who were 10-15 years older, suggesting that younger diabetic patients may require additional tools for risk stratification.

###

"Stress Myocardial Perfusion PET Provides Incremental Risk Prediction in Patients with and Patients without Diabetes." Collaborating with Dr. Skali were Marcelo F. Di Carli, M.D., Ron Blankstein, M.D., Benjamin J. Chow, M.D., Rob S. Beanlands, M.D., Daniel S. Berman, M.D., Guido Germano, Ph.D., James K. Min, M.D., Michael Merhige, M.D., Brent Williams, Ph.D., Emir Veledar, Ph.D., Leslee J. Shaw, Ph.D., and Sharmila Dorbala M.D., M.P.H.

Radiology: Cardiothoracic Imaging is edited by Suhny Abbata, M.D., University of Texas Southwestern Medical Center, Dallas, and owned and published by the Radiological Society of North America, Inc. (<https://pubs.rsna.org/cardiothoracic-imaging>)

RSNA is an association of over 53,400 radiologists, radiation oncologists, medical physicists and related scientists, promoting excellence in patient care and health care delivery through education, research and technologic innovation. The Society is based in Oak Brook, Ill. ([RSNA.org](https://www.rsna.org))

For patient-friendly information on PET, visit [RadiologyInfo.org](https://radiologyinfo.org).