

820 Jorie Blvd Oak Brook, IL 60523 TEL 1-630-571-2670 FAX 1-630-571-7837 RSNA.org



RSNA Press Release

MRI Reveals Brain Damage in Obese Teens

Released: November 25, 2019

CHICAGO — Researchers using MRI have found signs of damage that may be related to inflammation in the brains of obese adolescents, according to a study being presented next week at the annual meeting of the Radiological Society of North America (RSNA).



At A Glance

- Signs of damage in the brains of obese adolescents seen on MRI may be related to inflammation.
- The pattern of damage correlated with some inflammatory markers like leptin, a hormone made by fat cells that helps regulate energy levels and fat stores.
- Brain changes found in obese adolescents related to important regions responsible for control of appetite, emotions and cognitive functions.

Pamela Bertolazzi, Ph.D.

Obesity in young people has become a significant public health problem. In the U.S., the percentage of children and adolescents affected by obesity has more than tripled since the 1970s, according to the Centers for Disease Control and Prevention. Data from the World Health Organization indicates that the number of overweight or obese infants and young children ages five years or younger increased from 32 million globally in 1990 to 41 million in 2016.

While obesity is primarily associated with weight gain, recent evidence suggests that the

disease triggers inflammation in the nervous system that could damage important regions of the brain. Developments in MRI like diffusion tensor imaging (DTI), a technique that tracks the diffusion of water along the brain's signal-carrying white matter tracts, have enabled researchers to study this damage directly.

For the new study, researchers compared DTI results in 59 obese adolescents and 61 healthy adolescents, ages 12 to 16 years. From DTI, the researchers derived a measure called fractional anisotropy (FA), which correlates with the condition of the brain's white matter. A reduction in FA is indicative of increasing damage in the white matter.

The results showed a reduction of FA values in the obese adolescents in regions located in the corpus callosum, a bundle of nerve fibers that connects the left and right hemispheres of the brain. Decrease of FA was also found in the middle orbitofrontal gyrus, a brain region related to emotional control and the reward circuit. None of the brain regions in obese patients had increased FA.

"Brain changes found in obese adolescents related to important regions responsible for control of appetite, emotions and cognitive functions," said study co-author Pamela Bertolazzi, a biomedical scientist and Ph.D. student from the University of São Paulo in Brazil.

This pattern of damage correlated with some inflammatory markers like leptin, a hormone made by fat cells that helps regulate energy levels and fat stores. In some obese people, the brain does not respond to leptin, causing them to keep eating despite adequate or excessive fat stores. This condition, known as leptin resistance, makes the fat cells produce even more leptin.

Worsening condition of the white matter was also associated with levels of insulin, a hormone produced in the pancreas that helps regulate blood sugar levels. Obese people often suffer from insulin resistance, a state in which the body is resistant to the effects of the hormone.

"Our maps showed a positive correlation between brain changes and hormones such as leptin and insulin," Dr. Bertolazzi said. "Furthermore, we found a positive association with inflammatory markers, which leads us to believe in a process of neuroinflammation besides insulin and leptin resistance."

Dr. Bertolazzi noted that additional studies are needed to determine if this inflammation in young people with obesity is a consequence of the structural changes in the brain.

"In the future, we would like to repeat brain MRI in these adolescents after multi-professional treatment for weight loss to assess if the brain changes are reversible or not," she added.

Co-authors are Ricardo Uchida, D.Sc., Fabio L. Duran, D.Sc., Thaysa Neves, M.Sc., Elie Calfat, M.Sc., Naomi Costa, Estefania S. Fernandez, JoAnna D. Lima, Daniel A. Vasques, M.Sc., Cristiane Kochi, D.Sc., Marília Seelaender, D.Sc., Victor H. Otani, D.Sc., and Thais Z. Otani, D.Sc.

###

Note: Copies of RSNA 2019 news releases and electronic images will be available online at

<u>RSNA.org/press19</u> beginning Monday, Nov. 25.

RSNA is an association of over 53,400 radiologists, radiation oncologists, medical physicists and related scientists, promoting excellence in patient care and health care delivery through education, research and technologic innovation. The Society is based in Oak Brook, Ill. (*RSNA.org*)

Editor's note: The data in these releases may differ from those in the published abstract and those actually presented at the meeting, as researchers continue to update their data right up until the meeting. To ensure you are using the most up-to-date information, please call the RSNA Newsroom at 1-312-791-6610.

For patient-friendly information on brain MRI, visit <u>RadiologyInfo.org</u>.