

---

## RSNA Press Release

### Medical Radiation Exposure Fell in the U.S. from 2006 to 2016

Released: March 17, 2020

OAK BROOK, Ill. — Medical radiation exposure to patients in the U.S. fell by 20% between 2006 and 2016, reversing a quarter century-long trend of increasing exposure, according to a study appearing in the journal *Radiology*.

The use of medical imaging has grown rapidly in recent decades, raising concerns about the exposure of patients to ionizing radiation. A landmark report published in 2008 found that per capita radiation exposure in the U.S. increased six-fold between 1980 and 2006.

“The radiation dose to the U.S. population went up dramatically because of medical exposure, mostly from CT scanning and nuclear medicine, and that woke everybody up to the problem,” said study senior author Fred A. Mettler Jr., M.D., radiologist from the Department of Radiology at the University of New Mexico in Albuquerque.

In the wake of the report, medical societies and organizations enacted initiatives to increase awareness of exposure while equipment manufacturers developed more refined dose modulation technology. The effects of these and other efforts in the years since 2006 have remained largely unknown.

#### At A Glance

- Between 2006 and 2016, medical radiation exposure to patients in the U.S. fell by 20%.
- A key factor in the reduction was a substantial decrease in the number of nuclear medicine procedures, from 17 million in 2006 to 13.5 million in 2016.
- Researchers credit campaigns such as Image Wisely and Image Gently for optimizing dose and reducing unnecessary imaging examinations.



*Mahadevappa Mahesh, M.S., Ph.D.*

Armed with a grant from the Centers for Disease Control and Prevention, Dr. Mettler assembled a group of experts in medical imaging and physics and set out to determine the change in per capita radiation exposure in the U.S. from 2006 to 2016.

The results showed that the number of diagnostic and interventional radiology examinations performed remained largely unchanged over the 10-year period, even though the U.S. population increased by about 23 million. Estimated annual individual dose from diagnostic and interventional medical procedures fell from 2.9 millisieverts (mSv) in 2006 to 2.3 mSv in 2016, a decrease of approximately 20%.

“The overall trend stabilized, and the total dose to the U.S. population dropped a bit,” Dr. Mettler said.

A key factor in the reduction was a substantial decrease in the number of nuclear medicine procedures, from 17 million in 2006 to 13.5 million in 2016. The decline was particularly notable in cardiology after a cut in Medicare reimbursement drove many cardiologists away from nuclear medicine-based procedures to stress echocardiography, a test that relies on ultrasound instead of ionizing radiation.

“Nuclear medicine basically fell off a cliff, possibly due to cardiologists discovering that reimbursement was way down and that, for cardiac ischemia and other indications, stress echocardiography is essentially as accurate as the nuclear medicine procedures,” Dr. Mettler said.

CT scans, a major driver of medical radiation exposure, increased from 67 million to 84 million scans over the 10-year period. However, the average individual effective dose from CT procedures dropped by 6%, thanks to several factors, according to study co-author Mahadevappa Mahesh, M.S., Ph.D., professor of radiology and cardiology at Johns Hopkins University School of Medicine in Baltimore.

“One important factor is the dose modulation techniques available on most CT scanners in

the country,” he said. “The second factor is that overall CT detectors are becoming very efficient in the sense that they can utilize less radiation to create the same quality images.”

Dr. Mahesh also credited campaigns such as Image Wisely, a joint effort of the Radiological Society of North America, the American College of Radiology, American Society of Radiologic Technologists and American Association of Physicists in Medicine, aimed at optimizing dose and reducing unnecessary imaging examinations in the adult population, and Image Gently, which does the same in the pediatric population. He also pointed to accreditation requirements that are contributing towards dose stabilization.

Now that U.S. medical radiation exposure levels appear to have stabilized, the researchers stressed the importance of sustained vigilance to keep the trend from reversing.

“We don’t want people to become complacent,” Dr. Mahesh said. “Things are generally going in a good direction, but we need to continue on that path.”

Dr. Mahesh and colleagues plan to publish a report next year on trends in medical radiation exposure worldwide.

###

“Patient Exposure from Radiologic and Nuclear Medicine Procedures in the United States: Procedure Volume and Effective Dose for the Period 2006-2016.” Collaborating with Drs. Mettler and Mahesh were Mythreyi Bhargavan-Chatfield, Ph.D., Charles E. Chambers, M.D., Jennifer G. Elee, B.S., Donald P. Frush, M.D., Donald L. Miller, M.D., Henry D. Royal, M.D., Michael T. Milano, M.D., David C. Spelic, Ph.D., Armin J. Ansari, Ph.D., Wesley E. Bolch, Ph.D., Gary M. Guebert, D.C., Robert H. Sherrier, M.D., James M. Smith, Ph.D., and Richard J. Vetter, Ph.D.

*Radiology* is edited by David A. Bluemke, M.D., Ph.D., University of Wisconsin School of Medicine and Public Health, Madison, Wisconsin, and owned and published by the Radiological Society of North America, Inc. (<https://pubs.rsna.org/journal/radiology>)

RSNA is an association of radiologists, radiation oncologists, medical physicists and related scientists promoting excellence in patient care and health care delivery through education, research and technologic innovation. The Society is based in Oak Brook, Illinois. ([RSNA.org](https://www.rsna.org))

For patient-friendly information on radiation dose and patient safety, visit [RadiologyInfo.org](https://www.radiologyinfo.org).