
RSNA Press Release

Blood Flows Differently Through the Brains of Schizophrenic Patients

Released: May 26, 2010

Media Contacts:

RSNA Media Relations: 1-630-590-7762

Linda Brooks
1-630-590-7738
lb Brooks@rsna.org

Maureen Morley
1-630-590-7754
mmorley@rsna.org

OAK BROOK, Ill. — Researchers in Germany have used a magnetic resonance imaging (MRI) technique called continuous arterial spin labeling (CASL) to map cerebral blood flow patterns in schizophrenic patients quickly and without using radiation or contrast agents. Their findings appear in the online edition and July printed issue of the journal *Radiology*.

"Arterial spin labeling is a powerful technique that can help reduce the cost and complexity of examinations," said the study's lead author, Lukas Scheef, M.D., from the Department of Radiology at University of Bonn, Germany. "It can also be more readily repeated than methods that involve the use of contrast agents and radiotracers."

Schizophrenia is a chronic and severe brain disorder that affects approximately 2.4 million American adults, according to the National Institute of Mental Health. Symptoms can include hallucinations, delusions, disordered thinking, movement disorders, social withdrawal and cognitive deficits.

In the study, conducted at the University Hospital of Bonn in Germany, researchers used CASL MRI to compare cerebral blood flow in 11 non-medicated patients with schizophrenia and 25 healthy controls. The patient group included three women with a mean age of 36 years and eight men with a mean age of 32 years. The control group included 13 women (mean age, 29 years) and 12 men (mean age, 30 years).

The results revealed that compared to the healthy controls, the schizophrenic patients had extensive areas of hypoperfusion, or lower blood flow than normal in the frontal lobes and frontal cortex, anterior and medial cingulate gyri, and parietal lobes. These regions are associated with a number of higher cognitive functions including planning, decision making,

At A Glance

- Schizophrenic patients have reduced blood flow in the frontal lobes of the brain and increased blood flow in cerebellum, brainstem and thalamus.
- Continuous arterial spin labeling is a noninvasive technique used to measure the blood flow in the brain without the use of ionizing radiation.
- Schizophrenia affects approximately 2.4 million adults in the U.S.

judgment and impulse control.

Hyperperfusion, or increased blood flow, was observed in the cerebellum, brainstem and thalamus of the schizophrenic patients.

"Our CASL study revealed patterns of hypo- and hyperperfusion similar to the perfusion patterns observed in positron emission tomography (PET) and single photon emission computed tomography (SPECT) studies of schizophrenic patients," Dr. Scheef said.

Unlike PET and SPECT studies, CASL MR images can be quickly acquired without the use of ionizing radiation or contrast agents. In CASL MRI, arterial blood water is magnetically labeled in order to non-invasively measure cerebral blood flow.

"CASL MRI may allow researchers to gain a better understanding of schizophrenia," Dr. Scheef said. "In the long run, it may help to individualize and optimize treatment."

#

"Resting-State Perfusion in Non-Medicated Schizophrenic Patients: A Continuous Arterial Spin-Labeling 3.0-T MR Study." Collaborating with Dr. Scheef were Christoph Manka, M.D., Marcel Daamen, Kai-Uwe Kühn, M.D., Wolfgang Maier, M.D., Hans H. Schild, M.D., and Frank Jessen, M.D.

Radiology is edited by Herbert Y. Kressel, M.D., Harvard Medical School, Boston, Mass., and owned and published by the Radiological Society of North America, Inc. (<http://radiology.rsna.org/>)

RSNA is an association of more than 44,000 radiologists, radiation oncologists, medical physicists and related scientists committed to excellence in patient care through education and research. The Society is based in Oak Brook, Ill. (RSNA.org)

For patient-friendly information on MRI, visit RadiologyInfo.org.