

820 Jorie Blvd Oak Brook, IL 60523 TEL 1-630-571-2670 FAX 1-630-571-7837 RSNA.org



## **RSNA Press Release**

## Digital Mammography Cancer Detection Rates May Vary Significantly

Released: May 14, 2013

Media Contacts:

 RSNA Media Relations:
 1-630-590-7762

 Linda Brooks
 Maureen Morley

 1-630-590-7738
 1-630-590-7754

 Ibrooks@rsna.org
 mmorley@rsna.org

OAK BROOK, Ill. — Digital direct radiography (DR) is significantly more effective than computed radiography (CR) at detecting breast cancer, according to a new study published online in the journal *Radiology*.

The findings are the first to show a difference between the two types of digital imaging when compared with screen film mammography and suggest that women should be informed of the potential for lower cancer detection with CR, the researchers said.

Digital mammography, which takes an electronic image of the breast that can be stored and sent electronically, has supplanted screen film mammography in recent years.

DR and CR are the two types of digital mammography offered. DR is an online system in which the detector is an integral part of the mammographic unit, and the digital image can be read by the system in real time. CR is an offline system that relies on a cassette-based removable detector. An external reading device is used to generate the digital image.

While numerous studies have been published comparing digital mammography with screen film mammography, less is known about the comparative effectiveness of DR and CR, according to Anna M. Chiarelli, Ph.D., senior scientist in Prevention and Cancer Control at Cancer Care Ontario in Toronto, Canada.

"Digital mammography was implemented progressively in Ontario starting in 2006, and since that time there's been no evidence in the literature to suggest that CR did not



## At A Glance

- A study of nearly 700,000 women found that digital direct radiography is significantly more effective than computed radiography at detecting breast cancer.
- Researchers suggest that women should be informed of the potential for lower cancer detection with computed radiography.
- Computed radiography was 21 percent less effective than digital direct radiography, which could result in about 10 fewer cancers detected per 10,000 women screened.

perform as well as DR," she said.

For the new study, Dr. Chiarelli and colleagues drew upon information from the Ontario Breast Screening Program (OBSP), initiated in 1990 to deliver breast screening to women between the ages of 50 and 74. The researchers identified three groups of women 50 to 74 years old who were screened between Jan. 1, 2008, and Dec. 31, 2009. A total of 403,688 women were screened by screen film



Anna M. Chiarelli, Ph.D.

mammography, while 220,520 had DR and 64,210 underwent CR. The women were followed for 12 months after screening.

DR detected 4.9 cancers per 1,000 mammograms, a figure almost identical to screen film mammography's rate of 4.8 cancers per 1,000 mammograms. However, CR's detection rate of 3.4 cancers per 1,000 mammograms was significantly lower.

"CR was 21 percent less effective than DR," Dr. Chiarelli said. "This could result in about 10 fewer cancers detected per 10,000 women screened."

The reason for the lower detection rates likely can be attributed to technical factors, Dr. Chiarelli added.

"There may be several technical reasons reported by others for the lower effectiveness of CR, including loss of spatial resolution, or sharpness, and increased image noise, or granularity," she said.

Dr. Chiarelli said the new results carry more weight than those of previous studies because of the larger number of women screened and the reliance on concurrent cohorts, or different groups followed over the same time period.

###

"Digital Compared with Screen-Film Mammography: Performance Measures in Concurrent Cohorts within an Organized Breast Screening Program." Collaborating with Dr. Chiarelli were Sarah A. Edwards, M.H.Sc., Maegan V. Prummel, M.P.H., Derek Muradali, M.D., Vicky Majpruz, M.Sc., Susan J. Done, M.B., B.Chir., Patrick Brown, Ph.D., Rene S. Shumak, M.D., and Martin J. Yaffe, Ph.D. The study was funded by the Canadian Institutes of Health Research.

*Radiology* is edited by Herbert Y. Kressel, M.D., Harvard Medical School, Boston, Mass., and owned and published by the Radiological Society of North America, Inc. (<a href="http://radiology.rsna.org/">http://radiology.rsna.org/</a>)

RSNA is an association of more than 51,000 radiologists, radiation oncologists, medical physicists and related scientists, promoting excellence in patient care and health care delivery through education, research and technologic innovation. The Society is based in Oak Brook, Ill. (RSNA.org)

For patient-friendly information on digital mammography, visit RadiologyInfo.org.