

820 Jorie Blvd Oak Brook, IL 60523 TEL 1-630-571-2670 FAX 1-630-571-7837 RSNA.org



## **RSNA Press Release**

## **Breast Tomosynthesis Increases Cancer Detection and Reduces Recall Rates**

Released: December 3, 2013

Media Contacts: RSNA Newsroom 1-312-949-3233
Before 11/30/13 RSNA Media 1-630-590-7762

or after 12/5/13: Relations:

Linda Brooks Maureen Morley
1-630-590-7738 1-630-590-7754
Ibrooks@rsna.org mmorley@rsna.org

CHICAGO – Researchers have found that digital breast tomosynthesis (DBT) led to reduced recall rates and an increase in cancer detection in a large breast cancer screening program. The results of this study were presented today at the annual meeting of the Radiological Society of North America (RSNA).

Digital mammography is the gold standard for breast cancer screening, but may yield suspicious findings that turn out not to be cancer. These false-positive findings are associated with a higher recall rate, or the rate at which women are called back for additional imaging or biopsy.

Digital breast tomosynthesis has shown promise at reducing recall rates in all groups of patients, including younger women and women with dense breast tissue. Tomosynthesis is similar to

mammography in



Emily F. Conant, M.D.

 Researchers found that digital breast tomosynthesis (DBT) led to reduced recall rates

and an increase in cancer

At A Glance

detection.

- The research team compared the results from 15,633 women imaged with DBT to those of 10,753 patients imaged with digital mammography.
- The average recall rate decreased from 10.40 percent to 8.78 percent with DBT, and cancer detection rates increased from 4.28 to 5.25 (per 1,000 patients) with DBT.

that it relies on ionizing radiation to generate images of the breast. However, unlike conventional mammography, tomosynthesis allows for three-dimensional (3-D) reconstruction of the breast tissue, which can then be viewed as sequential slices through

the breast.

Because DBT technology is relatively new, it is typically used only as a supplemental screening tool, but since October 2011, every patient screened for breast cancer at Hospital of the University of Pennsylvania (HUP) in Philadelphia has been screened using DBT, according to Emily F. Conant, M.D., chief of breast imaging at HUP and the study's lead author.

"We have used DBT on all of our breast screening patients," Dr. Conant said. "Every patient has had it—we have not selected patients because of their risk or breast density or if they were willing to pay extra. We did not charge extra and were able to provide all of our women with this new technology."

For the study, Dr. Conant and colleagues compared imaging results from 15,633 women who underwent DBT at HUP beginning in 2011 to those of 10,753 patients imaged with digital mammography the prior year. Six radiologists trained in DBT interpretation reviewed the images.

The researchers found that, compared to digital mammography, the average recall rate using DBT decreased from 10.40 percent to 8.78 percent, and the cancer detection rate increased from 4.28 to 5.25 (per 1,000 patients). The overall positive predictive value—the proportion of positive screening mammograms from which cancer was diagnosed—increased from 4.1 percent to 6.0 percent with DBT.

"Our study showed that we reduced our callback rate and increased our cancer detection rate," Dr. Conant said. "The degree to which these rates were affected varied by radiologist. But importantly, the ratio of callback to cancer detection rate improved significantly for our radiologists."

Dr. Conant notes that tomosynthesis is an evolving platform, and researchers are already seeing a significant improvement in important screening outcomes.

"It's the most exciting improvement to mammography that I have seen in my career, even more important than the conversion from film-screen mammography to digital mammography," she said. "The coming years will be very exciting, as we see further improvements in this technology."

Co-authors are Nandita Mitra, Ph.D., Anne Marie McCarthy, Ph.D., Despina Konto, Ph.D., Susan G. Roth, M.D., Susan P. Weinstein, M.D., Marie Synnestvedt, Ph.D., Mathew Thomas, B.S., and Fei Wan, Ph.D.

###

Note: Copies of other RSNA 2013 news releases and electronic images will be available online at RSNA.org/press13 beginning Monday, Dec. 2.

RSNA is an association of more than 53,000 radiologists, radiation oncologists, medical physicists and related scientists, promoting excellence in patient care and health care delivery through education, research and technologic innovation. The Society is based in Oak Brook, Ill. (RSNA.org)

Editor's note: The data in these releases may differ from those in the published abstract and those actually presented at the meeting, as researchers continue to update their data right up until the meeting. To ensure you are using the most up-to-date information, please call the RSNA Newsroom at 1-312-949-3233.

For patient-friendly information on breast cancer screening, visit <u>RadiologyInfo.org</u> .	