

RSNA Press Release

Breast Cancer Risk Related to Changes in Breast Density as Women Age

Released: December 3, 2013

Media Contacts: **RSNA Newsroom** **1-312-949-3233**
Before 11/30/13 **RSNA Media** **1-630-590-7762**
or after 12/5/13: **Relations:**

Linda Brooks Maureen Morley
1-630-590-7738 1-630-590-7754
lbrooks@rsna.org mmorley@rsna.org

CHICAGO – Automated breast density measurement is predictive of breast cancer risk in younger women, and that risk may be related to the rate at which breast density changes in some women as they age, according to research being presented today at the annual meeting of the Radiological Society of North America (RSNA).

Breast density, as determined by mammography, is already known to be a strong and independent risk factor for breast cancer. The American Cancer Society (ACS) considers women with extremely dense breasts to be at moderately increased risk of cancer and recommends they talk with their doctors about adding magnetic resonance imaging (MRI) screening to their yearly mammograms.

"Women under age 50 are most at risk from density-associated breast cancer, and breast cancer in younger women is frequently of a more aggressive type, with larger tumors and a higher risk of recurrence," said the study's senior author, Nicholas Perry, M.B.B.S., FRCS, FRCR, director at the London Breast Institute in London, U.K.

For the new study, Dr. Perry and colleagues compared breast density and cancer risk between younger and older women and analyzed how the risk relates to changes in breast density over time. The study group included 282 breast cancer cases and 317 healthy control participants who underwent full-field digital mammography, with breast density measured separately

At A Glance

- The rate at which breast density changes in some women as they age may affect their breast cancer risk.
- Researchers compared breast density and cancer risk between younger and older women and analyzed how the risk relates to changes in breast density over time.
- Breast density, as determined by mammography, is known to be a strong and independent risk factor for breast cancer.



*Nicholas Perry, M.B.B.S., FRCS,
FRCR*

using an automated volumetric system.

"In general, we refer to breast density as being determined by mammographic appearance, and that has, by and large, in the past been done by visual estimation by the radiologist—in other words, subjective and qualitative," Dr. Perry said. "The automated system we used in the study is an algorithm that can be automatically and easily applied to a digital mammogram, which allows an objective and, therefore, quantitative density measurement that is reproducible."

Breast cancer patients showed higher mammographic density than healthy participants up to the age of 50. The healthy controls demonstrated a significant decline in density with age following a linear pattern, while there was considerably more variability in density regression among the breast cancer patients.

"The results are interesting, because there would appear to be some form of different biological density mechanism for normal breasts compared to breasts with cancer, and this appears to be most obvious for younger women," Dr. Perry said. "This is not likely to diminish the current ACS guidelines in any way, but it might add a new facet regarding the possibility of an early mammogram to establish an obvious risk factor, which may then lead to enhanced screening for those women with the densest breasts."

For instance, some women might undergo a modified exposure exam at age 35 to establish breast density levels, Dr. Perry noted. Those with denser breast tissue could then be followed more closely with mammography and additional imaging like MRI or ultrasound for earlier cancer detection and treatment.

"It has been estimated that about 40 percent of life years lost to breast cancer are from women under 50 diagnosed outside of screening programs," Dr. Perry said. "In my practice, which is largely composed of urban professional women, 40 percent of cancers year to year are diagnosed in women under 50, and 10 percent in women younger than 40."

Co-authors are Katja Pinker-Domenig, M.D., Kefah Mokbel, M.B.B.S., FRCS, Sue E. Milner, B.Sc., and Stephen W. Duffy, M.Sc.

###

Note: Copies of other RSNA 2013 news releases and electronic images will be available online at [RSNA.org/press13](https://www.rsna.org/press13) beginning Monday, Dec. 2.

RSNA is an association of more than 53,000 radiologists, radiation oncologists, medical physicists and related scientists, promoting excellence in patient care and health care delivery through education, research and technologic innovation. The Society is based in Oak Brook, Ill. ([RSNA.org](https://www.rsna.org))

Editor's note: The data in these releases may differ from those in the published abstract and those actually presented at the meeting, as researchers continue to update their data right up until the meeting. To ensure you are using the most up-to-date information, please call the RSNA Newsroom at 1-312-949-3233.

For patient-friendly information on mammography, visit [RadiologyInfo.org](https://www.RadiologyInfo.org).