

820 Jorie Blvd Oak Brook, IL 60523 TEL 1-630-571-2670 FAX 1-630-571-7837 RSNA.org



## **RSNA Press Release**

## CT Angiography Links Arterial Plaque with Diabetes, Blood Pressure, Cholesterol

Released: June 2, 2015

OAK BROOK, III. – Non-calcified arterial plaque is associated with diabetes, high systolic blood pressure and elevated "bad" cholesterol levels in asymptomatic individuals, according to a new



David A. Bluemke, M.D., Ph.D.

study published online in the journal Radiology.

Coronary artery disease (CAD) is the leading cause of death in men and women worldwide, accounting for 17 million deaths annually. Current treatment strategies focus on cardiovascular risk and serum

## At A Glance

- Coronary computed tomography angiography (CCTA) has emerged as a viable screening option for plaque, including non-calcified plaque.
- Non-calcified arterial plaque is associated with diabetes, high systolic blood pressure and elevated "bad" cholesterol levels in asymptomatic individuals.
- Coronary artery disease (CAD) is the leading cause of death in men and women worldwide.

cholesterol levels rather than direct assessment of extent of disease in the coronary arteries.

Plaque that forms in the arterial walls can restrict blood flow and, in some cases, rupture, leading to potentially fatal heart attacks. There is considerable evidence that calcified, or stable, plaque is less prone to rupture than non-calcified, or soft, plaque. Intravascular ultrasound can quantify non-calcified and calcified coronary artery plaque, but it is invasive and unsuitable for screening purposes. Coronary artery calcium (CAC) scoring with CT, a common noninvasive option, measures how much calcified plaque a person has, but it doesn't measure non-calcified plaque, and that's the component that tends to be dangerous. Despite treatment for hypercholesterolemia (high levels of cholesterol in the blood), CAC scores often paradoxically increase. Thus, researchers have searched for other plaque measures that can identify treatment response.

"Most information to date about coronary artery disease and cardiovascular risk factors in asymptomatic individuals has been derived from calcium scoring," said the study's lead author, David A. Bluemke, M.D., Ph.D., from the NIH Clinical Center. "We hypothesized that risk factors for the presence of non-calcified plaque may differ from those for calcified

plaque."

Coronary computed tomography angiography (CCTA) has emerged as a viable screening option for plaque, including non-calcified plaque. CCTA can capture the full anatomic map of the coronary arteries in a single heartbeat with low radiation dose and provide a complete picture of the total amount of plaque throughout the arteries of the heart.

For the study, Dr. Bluemke and colleagues used CCTA to assess the relationship between calcified and non-calcified coronary plaque burden in the coronary arteries and cardiovascular risk factors in low- to moderate-risk asymptomatic individuals.

The researchers recruited 202 asymptomatic men and women over age 55 who were eligible for statin therapy. CCTA was performed using a 320-detector row CT scanner and an intravenous contrast agent. Coronary wall thickness/plaque was evaluated, and analysis was performed to determine the relationship between risk factors and plaque.

Controlling for all risk factors, total coronary plaque index was greater in men than in women. Non-calcified plaque index was significantly associated with greater systolic blood pressure, diabetes, and elevated low-density lipoprotein (LDL) cholesterol level.

"These results highlight the potential of CCTA in quantifying plaque burden to assess progression or regression of coronary artery disease in low- to moderate-risk individuals," Dr. Bluemke said.

###

"Coronary Plaque Burden at Coronary CT Angiography in Asymptomatic Men and Women." Collaborating with Dr. Bluemke were Karen Rodriguez, B.A., Alan C. Kwan, B.A., Shenghan Lai, M.D., João A. C. Lima, M.D., Davis Vigneault, B.S., Veit Sandfort, M.D., Puskar Pattanayak, M.D., Mark A. Ahlman, M.D., Marissa Mallek, R.N., and Christopher T. Sibley, M.D.

*Radiology* is edited by Herbert Y. Kressel, M.D., Harvard Medical School, Boston, Mass., and owned and published by the Radiological Society of North America, Inc. (<u>http://Radiology.RSNA.org</u>)

RSNA is an association of more than 54,000 radiologists, radiation oncologists, medical physicists and related scientists promoting excellence in patient care and health care delivery through education, research and technologic innovation. The Society is based in Oak Brook, Ill. (RSNA.org)

For patient-friendly information on <u>CCTA</u>, visit <u>RadiologyInfo.org</u>.