

---

## RSNA Press Release

### Substance Abuse Reduces Brain Volume in Women but Not Men

Released: July 14, 2015

OAK BROOK, Ill. — Stimulant drug abuse has long-term effects on brain volume in women, according to a new study published online in the journal *Radiology*. Brain structures involved in reward, learning and executive

control showed vast changes even after a prolonged period of abstinence from drug use.

"We found that after an average of 13.5 months of abstinence, women who were previously dependent on stimulants had significantly less gray matter volume in several brain areas compared to healthy women," said the study's senior author, Jody Tanabe, M.D., professor of radiology, vice chair of Research, and Neuroradiology Section Chief at the University of Colorado Denver School of Medicine. "These brain areas are important for decision making, emotion, reward processing and habit formation."

For the study, Dr. Tanabe and colleagues sought to determine how the brains of people previously dependent on stimulants differed from the brains of healthy people. "We specifically wanted to determine how these brain effects differed by gender," Dr. Tanabe said.

The researchers analyzed structural brain magnetic resonance imaging (MRI) exams in 127 men and women, including 59 people (28 women and 31 men) who were previously dependent on cocaine, amphetamines, and/or methamphetamine for an average of 15.7 years, and 68 healthy people (28 women and 40 men) who were similar



*Michael Regner, M.D.*

#### At A Glance

- Women who were former stimulant drug abusers showed significant loss of gray matter volume in their brains.
- The women showed vast changes in brain structures that are important for decision making, emotion, reward processing and habit formation, compared to healthy women who were not prior substance abusers.
- Men who were former stimulant abusers demonstrated no significant brain differences compared to their healthy counterparts.



*Jody Tanabe, M.D.*

in age and gender. The MRI results showed that after an average of 13.5 months of abstinence, women who were previously dependent on stimulants had significantly less gray matter volume in frontal, limbic and temporal regions of the brain.

"While the women previously dependent on stimulants demonstrated widespread brain differences when compared to their healthy control counterparts, the men demonstrated no significant brain differences," Dr. Tanabe said.

The researchers also looked at how these brain volume differences were related to behaviors. They found that lower regional gray matter volumes correlated with behavioral tendencies to seek reward and novelty.

"Lower gray matter volumes in women who had been stimulant dependent were associated with more impulsivity, greater behavioral approach to reward, and also more severe drug use," Dr. Tanabe said. "In contrast, all men and healthy women did not show such correlations."

According to Dr. Tanabe, the results may provide a clue to the biological processes underlying the clinical course of stimulant abuse in men and women.

"Compared to men, women tend to begin cocaine or amphetamine use at an earlier age, show accelerated escalation of drug use, report more difficulty quitting and, upon seeking treatment, report using larger quantities of these drugs," she said. "We hope that our findings will lead to further investigation into gender differences in substance dependence and, thus, more effective treatments."

The study first author was Michael Regner, M.D., a radiology resident and PhD graduate student.

# # #

"Sex Differences in Gray Matter Changes and Brain-Behavior Relationships in Patients with Stimulant Dependence." Collaborating with Drs. Tanabe and Regner on this paper were Manish Dalwani, M.S., Dorothy Yamamoto, Ph.D., Robert I. Perry, M.D., Joseph T. Sakai, M.D., and Justin M. Honce, M.D.

*Radiology* is edited by Herbert Y. Kressel, M.D., Harvard Medical School, Boston, Mass., and owned and published by the Radiological Society of North America, Inc. (<http://radiology.rsna.org/>)

RSNA is an association of more than 54,000 radiologists, radiation oncologists, medical physicists and related scientists, promoting excellence in patient care and health care delivery through education, research and technologic innovation. The Society is based in Oak Brook, Ill. ([RSNA.org](http://RSNA.org))

For patient-friendly information on MRI of the brain, visit [RadiologyInfo.org](http://RadiologyInfo.org).