

820 Jorie Blvd Oak Brook, IL 60523 TEL 1-630-571-2670 FAX 1-630-571-7837 RSNA.org



RSNA Press Release

Imaging Links Structural Brain Changes and Cognitive Decline in Parkinson's

Released: December 7, 2016

OAK BROOK, Ill. — People with Parkinson's disease and cognitive impairment have disruptions in their brain networks that can be seen on a type of MRI, according to a study appearing online in the journal *Radiology*.

Parkinson's disease (PD) is a progressive disorder of the central nervous system characterized by tremors or trembling and stiffness in the limbs, impaired balance and coordination. It affects about 10



Massimo Filippi, M.D.

At A Glance

- Cognitive impairment in Parkinson's disease is likely the consequence of a disruption of complex structural brain networks.
- The results suggest that MRI may be helpful in monitoring Parkinson's disease and predicting the occurrence of cognitive complications.
- Parkinson's disease affects about 10 million people worldwide.

million people worldwide. As PD progresses, many patients develop mild cognitive impairment (MCI), a decline in cognitive abilities, including thinking, memory and language. MCI can be identified in approximately 25 percent of newly diagnosed PD patients, and patients with MCI progress to dementia more frequently than those with normal cognitive performance.

For the new study, lead investigator Massimo Filippi, M.D., from the Neuroimaging Research Unit at San Raffaele Scientific Institute in Milan, Italy, coauthors Federica Agosta, M.D., Ph.D., and Sebastiano Galantucci, M.D., and other colleagues used an MRI technique called diffusion tractography to look for differences in the neural networks of PD patients with and without MCI.

Increasingly, the human brain is understood as an integrated network, or connectome, that has both a structural and functional component. By applying an analytical tool called graph analysis to the imaging results, researchers can measure the relationships among highly connected and complex data like the network of connections in the human brain.

"Cognitive impairment in PD is one of the major non-motor complications of the disease, as

well as one of the major concerns of patients and caregivers at the time of diagnosis," Dr. Agosta said. "Study of the changes related to cognitive impairment in PD is imperative in order to be able to answer patients' questions and finally be able to predict the future development of this condition."

The study group was made up of 170 PD patients, including 54 with MCI and 116 without, and 41 healthy controls. Analysis of imaging results showed that only PD patients with MCI had significant alterations at the brain network level. Measurements of the movement and diffusion of water in the brain, an indicator of the condition of the brain's signal-carrying white matter, differentiated PD patients with MCI from healthy controls and non-MCI PD patients with a good accuracy. Researchers said the results show that cognitive impairment in PD is likely the consequence of a disruption of complex structural brain networks rather than degeneration of individual white matter bundles.

The results may offer markers to differentiate PD patients with and without cognitive deficits, according to Dr. Agosta.

"If confirmed and replicated by other studies, these results would suggest the use of MRI in PD to support the clinicians in monitoring the disease and predicting the occurrence of cognitive complications," she said.

The researchers have obtained resting state functional MRI data from the patients and plan to study the functional connectome alterations associated with cognitive impairment in PD and how structural and functional abnormalities are interrelated.

The study is part of an international collaboration between the San Raffaele Scientific Institute and the Clinic of Neurology at the University of Belgrade in Belgrade, Serbia, led by Vladimir Kostic, M.D.

###

"Structural Brain Connectome and Cognitive Impairment in Parkinson's Disease." Collaborating with Drs. Filippi, Agosta and Galantucci were Elka Stefanova, M.D., Silvia Basaia, M.Sc., Martijn P. van den Heuvel, Ph.D., Tanja Stojković, M.D., Elisa Canu, Ph.D., Iva Stanković, M.D., Vladana Spica, M.D., Massimiliano Copetti, Ph.D., Delia Gagliardi, M.D., and Vladimir S. Kostic, M.D.

Radiology is edited by Herbert Y. Kressel, M.D., Harvard Medical School, Boston, Mass., and owned and published by the Radiological Society of North America, Inc. (<u>http://radiology.rsna.org/</u>)

RSNA is an association of more than 54,000 radiologists, radiation oncologists, medical physicists and related scientists, promoting excellence in patient care and health care delivery through education, research and technologic innovation. The Society is based in Oak Brook, Ill. (*RSNA.org*)

For patient-friendly information on brain MRI, visit <u>RadiologyInfo.org</u>.