
RSNA Press Release

Radiologists Can Help Identify Victims of Domestic Violence

Released: February 5, 2019

OAK BROOK, Ill. — Radiologists may play a crucial role in identifying signs of intimate partner violence, a type of domestic violence, according to a study published in the journal *Radiology*. Radiologists can identify potential violence-related patterns of injury and work closely with referring providers to provide care for the victims.



Bharti Khurana, M.D.

At A Glance

- Medical images can offer early clues of intimate partner violence.
- Researchers found common injuries among victims of intimate partner violence.
- Radiologists have access to prior imaging records that can assist in revealing a history of abuse.

Intimate partner violence is a serious, preventable social and public health problem that affects millions of Americans. The term describes physical, sexual or psychological harm by a current or former partner or spouse.

"Radiology has made substantial contributions to the detection of non-accidental trauma in children, and identification of pediatric violence is an essential component of a radiologist's training, but little is known about the role of imaging in the detection of intimate partner violence," said Bharti Khurana, M.D., the study's principal investigator, a radiologist at Brigham and Women's Hospital and assistant professor of radiology at Harvard Medical School in Boston. "Medical images can offer early clues of IPV based on location and pattern of both new and old injuries."

For the study, electronic medical records from 185 patients referred by Brigham and Women's Hospital Emergency Department (ED) to the intimate partner violence support

program from January 2015 to October 2016 were reviewed and compared to an age- and sex-matched control group of 555 patients who were seen in the ED. Reports of all patient imaging exams performed within the preceding five years were also reviewed.

The researchers found that almost all of the intimate partner violence victims were women (96 percent). Victims had a mean age of 34.2 years. Thirty-seven percent were African-American and 28 percent were Hispanic. Victims of intimate partner violence, on average, underwent approximately four times more imaging exams in the preceding five years compared to the control group. Acute fractures and obstetric/gynecologic findings, such as pregnancy loss and intrauterine growth restriction, were associated with two to four times higher odds of domestic violence.



Elizabeth George, M.D.

"Our study suggests that these otherwise common findings could be indicative of intimate partner violence in the setting of specific socioeconomic factors identified in our study," said study first author Elizabeth George, M.D., chief resident in the Department of Radiology at Brigham and Women's Hospital and clinical fellow at Harvard Medical School.

The research team found that intimate partner violence victims were more likely to be homeless (28 percent), have a psychiatric history (64 percent) or have a substance use disorder (24 percent), compared to patients who were not victims. The team found that physical abuse was the most common form of intimate partner violence (84 percent), but that patients also complained of emotional and verbal abuse (44 percent). Sexual abuse was seen in 15 percent of victims.

"Radiologists are in a unique position to share objective findings unbiased by interactions with the victim and potential abuser," Dr. Khurana said. "Encouraged by the early results, we have now expanded the study to include colleagues from orthopedics, primary care, women's health, social work, legal, and two additional Boston hospitals: Massachusetts General Hospital and Boston Medical Center."

The ultimate goal of the research, according to Dr. Khurana, is to develop an integrated system that uses patterns derived from expert analysis of historical imaging and clinical data, detects and classifies injuries, and automatically alerts clinicians if a patient's injuries have low- or high-risk probability for being the result of intimate partner violence.

In the next phase of the study, the researchers plan to use machine learning to recognize signs of intimate partner violence and create an alert system for greater confidence and robustness in findings, empowering health care providers to open a dialogue with their patients.

"In addition, our multidisciplinary team will design conversational guides using medical images for training for social workers and clinicians to approach the patients who are identified as being at high risk for intimate partner violence and are not forthcoming," Dr. Khurana said.

#

"Radiologic Findings in Intimate Partner Violence." Collaborating with Drs. George and Khurana were Catherine H. Phillips, M.D., Nandish Shah, M.D., Annie Lewis-O'Connor, N.P., Ph.D., Bernard Rosner, Ph.D., and Hanni M. Stoklosa, M.D.

Radiology is edited by David A. Bluemke, M.D., Ph.D., University of Wisconsin School of Medicine and Public Health, Madison, Wis., and owned and published by the Radiological Society of North America, Inc. (<http://radiology.rsna.org/>)

RSNA is an association of over 53,400 radiologists, radiation oncologists, medical physicists and related scientists, promoting excellence in patient care and health care delivery through education, research and technologic innovation. The Society is based in Oak Brook, Ill. (RSNA.org)

For patient-friendly information on emergency radiology, visit RadiologyInfo.org.