

820 Jorie Blvd Oak Brook, IL 60523 TEL 1-630-571-2670 FAX 1-630-571-7837 **RSNA.org**



RSNA Press Release

Novel Technique Accurately Assesses Cardiovascular Risks

Released: February 12, 2019

OAK BROOK, Ill. — A new noninvasive technique for imaging the carotid artery offers advantages over other imaging methods and could provide an earlier, more accurate assessment of cardiovascular disease risk, according to a study published in the journal *Radiology*.



At A Glance

- A new technique for imaging carotid arteries may provide an earlier, more accurate assessment of cardiovascular disease risk.
- Volumetric multi-spectral optoacoustic tomography (vMSOT) investigates tissue at a molecular scale.
- vMSOT provides information about disease-related biomarkers early enough to provide better treatment options.

Daniel Razansky, Ph.D.

The carotid arteries are the blood vessels located on the left and right side of the neck that bring oxygenated blood to the head. Each artery bifurcates, or forks, in the neck into two branches that form the internal and external carotid arteries. Most ischemic strokes, or strokes related to a build-up of plaque in the arteries, are associated with carotid artery disease originating from the area where the arteries bifurcate.

Imaging techniques like ultrasound, CT and MRI are useful for revealing the extent of narrowing in the carotid arteries, but less helpful in determining the makeup of the plaque itself. This is a crucial limitation because plaque composition is associated with vulnerability to rupture, setting in motion the chain of events that leads to life-threatening strokes.

"Rapid characterization of tissue function and molecular composition is limited with these modalities, which commonly results in poor diagnostic accuracy and ineffective treatments," said study senior author Daniel Razansky, Ph.D., director of the Functional and Molecular Imaging Lab at the University of Zurich and the Swiss Federal Institute of Technology in

Zurich.

Dr. Razansky and colleagues studied a new technique for carotid artery assessment called volumetric multi-spectral optoacoustic tomography (vMSOT). As with ultrasound, vMSOT is performed with a handheld device that is moved against the neck. However, vMSOT employs the science of spectroscopy to investigate tissue at a molecular scale. This provides information about the artery that is not attainable with other methods. It also can detect lipids, the pigment melanin and other disease-related biomarkers early enough to provide better treatment options.

"Unlike most other clinical imaging modalities mainly looking at late-stage anatomical manifestations of diseases, vMSOT is capable of sensing specific molecules in tissues without administration of contrast agents," Dr. Razansky said. "In the case of carotid artery disease, assessment of the entire bifurcation area in real time and in 3-D is only possible with vMSOT."

The researchers performed vMSOT imaging on 16 healthy participants and compared results with those from conventional ultrasound. The vMSOT approach was able to noninvasively and instantaneously assess the entire bifurcation area of the carotid artery in three dimensions, thus making it less prone than ultrasound to motion-related, image-blurring artifacts. Researchers said the results point to the tremendous potential of the new approach in the clinic.

"The developed handheld vMSOT imaging approach holds promise for rapid volumetric assessment of the carotid artery and plaque vulnerability in an entirely noninvasive manner," Dr. Razansky said. "It also has the additional potential for label-free identification and assessment of clinically-relevant biomarkers of carotid artery disease, which helps with early and accurate diagnosis, timely treatment planning and monitoring."

In the future, vMSOT could be combined with ultrasound for a more comprehensive characterization of the carotid artery.

"Given its fast imaging performance, excellent molecular contrast, portability and affordability, I truly believe that vMSOT will soon be routinely used in the clinic," Dr. Razansky said. "One day, it may even become as popular as ultrasound."

###

"Real-time Volumetric Assessment of the Human Carotid Artery: Handheld Multispectral Optoacoustic Tomography." Collaborating with Dr. Razansky were Ivana Ivankovic, M.Sc., Elena Merčep, M.Sc., Claus-Georg Schmedt, M.D., and Xose Luís Deán-Ben, Ph.D.

Radiology is edited by David A. Bluemke, M.D., Ph.D., University of Wisconsin School of Medicine and Public Health, Madison, Wis., and owned and published by the Radiological Society of North America, Inc. (http://radiology.rsna.org/)

RSNA is an association of over 53,400 radiologists, radiation oncologists, medical physicists and related scientists, promoting excellence in patient care and health care delivery through education, research and technologic innovation. The Society is based in Oak Brook, Ill. (*RSNA.org*)

For patient-friendly information on cardiovascular imaging, visit <u>RadiologyInfo.org</u>.