
RSNA Press Release

Chest X-rays Show More Severe COVID-19 in Non-white Patients

Released: July 16, 2020

OAK BROOK, Ill. (July 16, 2020) — Racial/ethnic minority patients admitted to the hospital with COVID-19 infection are more likely to have more severe disease on chest X-rays than white/non-Hispanic patients, increasing the likelihood of adverse outcomes, such as intubation or death, according to a study published in the journal *Radiology*.

Emerging data show that racial/ethnic minorities have been disproportionately affected by COVID-19. Socioeconomic factors and pre-existing medical conditions like hypertension are likely contributing factors to this disparity. Furthermore, limited English proficiency may introduce additional linguistic and health literacy barriers to care, potentially resulting in delays seeking medical attention and greater severity of disease at the time of admission to the hospital with COVID-19 infection.

Radiologists from Massachusetts General Hospital (MGH) saw these disparities firsthand in April among patients admitted to the hospital with confirmed COVID-19 infection, and at one of the hospital's respiratory infection clinics in Chelsea, a city just north of Boston that is home to a predominantly Spanish-speaking Hispanic community. A significant proportion of the patients who visited the Chelsea clinic had COVID-19, and the level of disease the radiologists observed on chest imaging was markedly more severe than that seen at other respiratory infection clinics in Boston. These disparities were more evident among patients admitted to the hospital with confirmed COVID-19 infection.

“It got to the point where half of our patient population admitted with COVID-19 were underrepresented minorities,” said study coauthor Efren J. Flores, M.D., a radiologist at MGH.

Dr. Flores and colleagues set out to study these observed disparities in greater detail with an eye toward developing a better understanding of some of the factors involved. They hope to use this information to guide risk mitigation strategies and improve outcomes among

At A Glance

- Upon admission to the hospital, non-white patients with COVID-19 infection are more likely to have more severe disease on chest X-rays than white/non-Hispanic patients.
- Researchers looked at data from 326 patients hospitalized with confirmed COVID-19 infection between March 27 and April 10, 2020.
- Increased severity on chest X-rays increased the likelihood of adverse clinical outcomes, including admission to the intensive care unit, intubation and death.

racial/ethnic minority groups. They looked at data from 326 patients hospitalized with confirmed COVID-19 infection between March 27 and April 10, 2020. Analysis of chest X-ray results revealed that non-white patients had significantly more severe lung disease on admission than white/non-Hispanic patients. Increased disease severity on chest X-rays increased the likelihood of adverse clinical outcomes, including admission to the intensive care unit, intubation and death.



Efred J. Flores, M.D.

As expected, the increased severity of lung disease on chest X-rays among non-white patients correlated with a combination of factors, including delay in seeking care at the hospital, higher prevalence of pre-existing comorbidities and limited English proficiency.

“Limited English proficiency is an additional socioeconomic factor that really influences many aspects of access to care,” Dr. Flores said. “When we were first learning how the disease spreads, there was all this rapidly evolving information coming out that was not available in languages other than English, and that lag in availability of actionable health information for non-English speaking individuals was really critical for many patients trying to navigate a complex medical system with a disease from a virus that is so aggressive.”

The connection between limited English proficiency and disease severity underscores the importance of having multilingual, culturally tailored health information available, Dr. Flores said, especially as the number of infections climbs in different parts of the country.

Disparities in access to care and disease severity are not limited to linguistic barriers. Racial/ethnic non-white communities disproportionately experience lower socioeconomic status, adding another layer of complexity when accessing care. Living and working arrangements also likely played a role in the severity of COVID-19 among these patients, the study authors said. Racial and ethnic minority populations tend to live in multigenerational households and communities of higher population density, making social distancing difficult. In addition, they are more often employed in jobs not conducive to remote work with limited paid time off, thus increasing their exposure to COVID-19.

“Many of these patients delay their care because they’re considered essential workers and they don’t have a lot of sick leave, but also it’s difficult for them to leave because they are living on a weekly paycheck and have other dependents,” Dr. Flores said. “It wasn’t uncommon for us to go into the medical record when we were interpreting their exams and see that many of them worked at grocery stores or warehouses.”

The study findings highlight the important role radiologists play in providing earlier identification of higher-risk patients and developing multidisciplinary collaborations to help

address these disparities, according to Dr. Flores.

“Health equity is every medical specialty’s responsibility, but I believe radiology is uniquely positioned to take a bigger role not only in population health but in public health efforts,” he said. “Our ability to provide care in different settings is what allowed us to make the clinical observation that patients coming to this one particular clinic and those being admitted to the hospital were presenting with a higher rate of positive findings that were also more severe. That really offered a window into the disparities that were translating into greater disease severity and worse outcomes.”

Moving forward, the findings of this study could assist radiologists in the development of algorithms to identify vulnerable and at-risk populations. This could spur collaborations with other medical specialties, community stakeholders, and public health initiatives to enhance the effectiveness of public health interventions that increase access to care.

“We did this study not only to gain a better understanding of these emerging disparities, but also to discover how we can use this information to craft a better path towards equity together,” Dr. Flores said.

###

“Racial/Ethnic Disparities in Disease Severity on Admission Chest Radiographs among Patients Admitted with Confirmed COVID-19: A Retrospective Cohort Study.” Collaborating with Dr. Flores were Nicholas P. Joseph, B.A., Nicholas J. Reid, B.Sc., Avik Som, M.D., Ph.D., Matthew D. Li, M.D., Emily P. Hyle, M.D., Caitlin M. Dugdale, M.D., Min Lang, M.D., Joseph R. Betancourt, M.D., M.P.H., Francis Deng, M.D., Dexter P. Mendoza, M.D., Brent P. Little, M.D., and Anand K. Narayan, M.D., Ph.D.

Radiology is edited by David A. Bluemke, M.D., Ph.D., University of Wisconsin School of Medicine and Public Health, Madison, Wisconsin, and owned and published by the Radiological Society of North America, Inc. (<https://pubs.rsna.org/journal/radiology>)

RSNA is an association of radiologists, radiation oncologists, medical physicists and related scientists promoting excellence in patient care and health care delivery through education, research and technologic innovation. The Society is based in Oak Brook, Illinois. ([RSNA.org](https://www.rsna.org))

For patient-friendly information on chest X-ray, visit [RadiologyInfo.org](https://www.rsna.org/radiologyinfo).