
RSNA Press Release

New Research May Lead to Improved Diagnosis of Autism

Released: May 31, 2011

Media Contacts:

RSNA Media Relations: 1-630-590-7762

Linda Brooks
1-630-590-7738
lbrooks@rsna.org

Maureen Morley
1-630-590-7754
mmorley@rsna.org

OAK BROOK, Ill. — Functional magnetic resonance imaging (fMRI) may provide an early and objective indicator of autism, according to researchers at Columbia University in New York City, who used the technique to document language impairment in autistic children. Results of their study appear online and in the August issue of *Radiology*.

Autism is a spectrum disorder characterized by repetitive behaviors and impaired language, communication and social interactions. According to the Centers for Disease Control and Prevention, it is estimated that as many as one in every 110 children is affected by autism.

"With the extraordinarily high prevalence of autism, you would think there would be an objective diagnosis for the disorder," said Joy Hirsch, Ph.D., a professor at Columbia University Medical Center and director of the Functional MRI Laboratory. "However, the diagnosis of autism currently remains limited to parent and clinician observation of missed developmental milestones."

In the study, researchers performed fMRI exams on 15 control children (mean age: 12.1) and 12 language-impaired and age-matched autistic children (mean age: 12.4). Using fMRI, the researchers were able to measure neural activity in working brain tissues, while the children listened to recordings of their parents talking to them.

Activation levels during passive stimulation were measured within two regions of the brain: the primary auditory cortex (A1) and superior temporal gyrus (STG), a region associated with sentence comprehension. Brain activation maps for each patient were then computed using statistical linear modeling.

At A Glance

- Using fMRI and passive speech stimulation, researchers identified different patterns of brain activation in children with autism.
- In the brain region associated with sentence comprehension, the autistic patients differed from the control patients.
- As many as one in every 110 children is affected by autism.



Joy Hirsch, Ph.D.

Activity in the A1 region of the brain did not differ between autistic and control patients. However, activation within the STG was greater for control children relative to autistic patients.

"These findings first tell us that the autistic children in our study appeared normal with respect to the primary auditory system," Dr. Hirsch said. "But it appears that the STG in the autistic brains was not as sensitive to the language narratives as was the STG in the brains of the typical children."

An additional 27 autistic children undergoing routine MRI exams with sedation were also included in the study. Using a similar analysis of sedation-adjusted values from the control group, the researchers identified 26 of 27 (96 percent) sedated autistic patients with autism.

"This study suggests that fMRI acquired during listening to a language narrative can be used to distinguish children with autism from those without," Dr. Hirsch said. "Based on these initial findings, future studies using these or similar fMRI methods may result in an early and objective imaging indicator for autism."

Some children with autism spectrum disorders can benefit from intensive behavior therapy, but early intervention is key.

"The need for an early, objective diagnosis is enormous," Dr. Hirsch said.

###

"Speech Stimulation during Functional MR Imaging as a Potential Indicator of Autism." Collaborating with Dr. Hirsch were Grace Lai, Ph.D., Harry D. Schneider, M.D., and Johanna C. Schwarzenberger, M.D.

Radiology is edited by Herbert Y. Kressel, M.D., Harvard Medical School, Boston, Mass., and owned and published by the Radiological Society of North America, Inc. (<http://radiology.rsna.org/>)

RSNA is an association of more than 46,000 radiologists, radiation oncologists, medical physicists and related scientists committed to excellence in patient care through education and research. The Society is based in Oak Brook, Ill. (RSNA.org)

For patient-friendly information on fMRI, visit RadiologyInfo.org.