

---

## RSNA Press Release

### Brain Connectivity Disrupted in Patients with Post-Concussive Syndrome

Released: July 20, 2011

Media Contacts:

**RSNA Media Relations: 1-630-590-7762**

Linda Brooks  
1-630-590-7738  
[lbrooks@rsna.org](mailto:lbrooks@rsna.org)

Maureen Morley  
1-630-590-7754  
[mmorley@rsna.org](mailto:mmorley@rsna.org)

OAK BROOK, Ill. — A new study has found that patients with mild traumatic brain injury (MTBI) exhibit abnormal functional connectivity in the thalamus, a centrally located relay station for transmitting information throughout the brain. The results of the study appear online in the journal *Radiology*.


Using resting-state functional MRI, we found increased functional connectivity of thalamocortical networks in patients following MTBI, due to the subtle injury of the thalamus," said study co-author Yulin Ge, M.D., associate professor in the Department of Radiology at NYU Langone Medical Center. "These findings hold promise for better elucidating the underlying cause of a variety of post-traumatic symptoms that are difficult to spot and characterize using conventional imaging methods."

According to the Centers for Disease Control and Prevention, each year in the U.S. 1.5 million people sustain traumatic brain injuries, resulting from sudden trauma to the brain. MTBI, or concussion, accounts for at least 75 percent of all traumatic brain injuries. Following a concussion, some patients experience a brief loss of consciousness. Other symptoms include headache, dizziness, memory loss, attention deficit, depression and anxiety. Some of these conditions may persist for months or even years. Typically in patients with MTBI, there are no structural abnormalities visible on the brain, so researchers have begun using specialized imaging exams to detect abnormalities in how the brain functions.

Comparing levels of activity among different groups of brain cells helps identify which

#### At A Glance

- Connectivity between brain networks may be disrupted in patients with MTBI, commonly known as concussion.
- MTBI accounts for at least 75% of all traumatic brain injuries.
- This is the first study to use resting-state fMRI to look at thalamus network disruption after concussion.



*Yulin Ge, M.D.*

brain networks are communicating with one another. Some brain networks, known as resting state networks (RSNs) and baseline brain activities can be detected when the brain is at rest. These networks include the parts of the brain associated with working memory.

"The RSNs have great potential for studying thalamic dysfunction in several clinical disorders including traumatic brain injury," Dr. Ge said.

Resting-state functional MRI (RS-fMRI) has rapidly emerged as a novel informative tool for investigating brain connectivity between regions that are functionally linked. RS-fMRI provides insight into functional activity and communication between brain regions, which play key roles in cognitive performance.

"The disruption of such functional properties is better characterized by RS-fMRI than by conventional diagnostic tools," Dr. Ge said.

Dr. Ge and colleagues used RS-fMRI to study the brain activity of 24 patients with MTBI and 17 healthy control patients. A normal pattern of thalamic RSNs with relatively symmetric and restrictive connectivity was demonstrated in the healthy control group. In the patients with MTBI, this pattern was disrupted, with significantly increased thalamic RSNs and decreased symmetry. These findings correlated with clinical symptoms and diminished neurocognitive functions in the patients with MTBI.

"The thalamic functional networks have multiple functions, including sensory information process and relay, consciousness, cognition, and sleep and wakefulness regulation," Dr. Ge said. "The disruption of thalamic RSNs may result in a burning or aching sensation, accompanied by mood swings and sleep disorders, and can contribute to certain psychotic, affective, obsessive-compulsive, anxiety and impulse control disorders. These symptoms are commonly seen in MTBI patients with post-concussive syndrome."

Because the causes of post-concussive syndrome are poorly understood, there is currently no treatment. But, according to Dr. Ge, the results of this study have implications for a new therapeutic strategy, based on sound understanding of the underlying mechanisms of thalamocortical disruption and post-concussive syndrome.

###

"Thalamic Resting-State Functional Networks: Disruption in Patients with Mild Traumatic Brain Injury." Collaborating with Dr. Ge on this paper were Lin Tang, Ph.D., Daniel K. Sodickson, M.D., Ph.D., Laura Miles, Ph.D., Yongxia Zhou, Ph.D., Joseph Reaume, B.S.R.T., and Robert I. Grossman, M.D.

*Radiology* is edited by Herbert Y. Kressel, M.D., Harvard Medical School, Boston, Mass., and owned and published by the Radiological Society of North America, Inc. (<http://radiology.rsna.org/>)

RSNA is an association of more than 46,000 radiologists, radiation oncologists, medical physicists and related scientists committed to excellence in patient care through education and research. The Society is based in Oak Brook, Ill. ([RSNA.org](http://RSNA.org))

For patient-friendly information on [Functional Magnetic Resonance Imaging \(fMRI\)](#), visit [RadiologyInfo.org](http://RadiologyInfo.org).