
RSNA Press Release

Novel MRI Technique Could Reduce Breast Biopsies

Released: October 2, 2012

Media Contacts:

RSNA Media Relations: 1-630-590-7762

Linda Brooks
1-630-590-7738
lbrosks@rsna.org

Maureen Morley
1-630-590-7754
mmorley@rsna.org

OAK BROOK, Ill. — Water diffusion measurements with MRI could decrease false-positive breast cancer results and reduce preventable biopsies, according to a new study published online in the journal *Radiology*. Researchers said the technique also could improve patient management by differentiating high-risk lesions requiring additional workup from other non-malignant subtypes.

Dynamic contrast-enhanced MRI (DCE-MRI) has emerged in recent years as a useful tool in breast cancer detection and staging. One of its primary limitations is a substantial number of false-positive findings that require biopsies.

"Many benign lesions demonstrate enhancement on DCE-MRI," said Savannah C. Partridge, Ph.D., research associate professor at the University of Washington, Seattle Cancer Care Alliance. "We need another means for differentiating benign lesions from malignancies."

One possible solution is diffusion-weighted imaging (DWI), an MRI technique that calculates the apparent diffusion coefficient (ADC)—a measure of how water moves through tissue.

"DWI has been used mostly in neurological applications, but it's been studied more recently in breast imaging," Dr. Partridge said. "It only adds a couple of minutes to the MRI exam and does not require additional contrast or any extra hardware."

Research has shown that DWI is a promising tool for distinguishing between benign and

At A Glance

- Water diffusion measurements with MRI could decrease false-positive breast cancer results and reduce preventable biopsies.
- Research has shown that diffusion-weighted imaging is a promising tool for distinguishing between benign and malignant breast lesions.
- Diffusion-weighted imaging successfully characterized 46 percent of non-malignant breast lesions identified as false-positive findings on DCE-MRI.



Savannah C. Partridge, Ph.D.

malignant breast lesions. Normal breast tissue has a high ADC because water moves through it relatively freely, while most cancers have a lower ADC because their cells are more tightly packed and restrict water motion. However, significant overlap exists between the ADC values of non-malignant lesions and breast malignancies, and little is known about the ADC values of specific subtypes of non-malignant lesions.

For the new study, Dr. Partridge and colleagues evaluated the DWI characteristics of non-malignant lesions in 165 women. Based on ADC values above a previously determined diagnostic threshold, DWI successfully characterized 46 percent of non-malignant breast lesions identified as false-positive findings on DCE-MRI as benign.

"We were excited to see the number of false positives that could be reduced through this approach," Dr. Partridge said. "DWI gives us extra microstructural information to distinguish among lesions. We can use ADC values to draw a cutoff above which we might not need to do a biopsy."

The research team is planning a multicenter trial to validate the findings and determine how to best to incorporate ADC measures into clinical breast MRI interpretations.

"We are very motivated to translate this promising technology to a clinically useful breast imaging tool," Dr. Partridge said.

#

"Nonmalignant Breast Lesions: ADCs of Benign and High-Risk Subtypes Assessed as False-Positive at Dynamic Enhanced MR Imaging." Collaborating with Dr. Partridge were Sana Parsian, M.D., Habib Rahbar, M.D., Kimberly H. Allison, M.D., Wendy D. DeMartini, M.D., Matthew L. Olson, M.S., and Constance D. Lehman, M.D., Ph.D.

Radiology is edited by Herbert Y. Kressel, M.D., Harvard Medical School, Boston, Mass., and owned and published by the Radiological Society of North America, Inc. (<http://radiology.rsna.org/>)

RSNA is an association of more than 50,000 radiologists, radiation oncologists, medical physicists and related scientists promoting excellence in patient care and health care delivery through education, research and technologic innovation. The Society is based in Oak Brook, Ill. (RSNA.org)

For patient-friendly information on [breast MRI](#), visit RadiologyInfo.org.