

---

## RSNA Press Release

### MRI Technique Reveals Low Brain Iron in ADHD Patients

Released: December 2, 2013

**Media Contacts:**    **RSNA Newsroom**    **1-312-949-3233**  
**Before 11/30/13**    **RSNA Media**    **1-630-590-7762**  
**or after 12/5/13:**    **Relations:**  
  
Linda Brooks                      Maureen Morley  
1-630-590-7738                      1-630-590-7754  
[lbrooks@rsna.org](mailto:lbrooks@rsna.org)                      [mmorley@rsna.org](mailto:mmorley@rsna.org)

CHICAGO—Magnetic resonance imaging (MRI) provides a noninvasive way to measure iron levels in the brains of people with attention deficit hyperactivity disorder (ADHD), according to a study being presented today at the annual meeting of the Radiological Society of North America (RSNA). Researchers said the method could help physicians and parents make better informed decisions about medication.

ADHD is a common disorder in children and adolescents that can continue into adulthood. Symptoms include hyperactivity and difficulty staying focused, paying attention and controlling behavior.

The American Psychiatric Association reports that ADHD affects 3 to 7 percent of school-age children.

Psychostimulant medications such as Ritalin are among the drugs commonly used to reduce ADHD symptoms. Psychostimulants affect levels of dopamine, a neurotransmitter in the brain associated with addiction.

"Studies show that psychostimulant drugs increase dopamine levels and help the kids that we suspect have lower dopamine levels," said Vitria Adisetiyo, Ph.D., postdoctoral research fellow at the Medical University of South Carolina in Charleston, S.C. "As brain iron is required for dopamine synthesis, assessment of iron levels with MRI may provide a noninvasive, indirect measure of dopamine."

Dr. Adisetiyo and colleagues explored this possibility by measuring brain iron in 22 children and adolescents with ADHD and 27 healthy control children and adolescents using an MRI technique called magnetic field correlation

#### At A Glance

- Researchers measured brain iron in children and adolescents with ADHD using an MRI technique called magnetic field correlation (MFC) imaging.
- Results showed that ADHD patients who had never been on medication had significantly lower MFC than the control group or ADHD patients who had been on psychostimulant medication.
- The MFC technique's ability to noninvasively detect low iron levels may help improve ADHD diagnosis and guide optimal treatment.



(MFC) imaging. The technique is relatively new, having been introduced in 2006 by study co-authors and faculty members Joseph A. Helpert, Ph.D., and Jens H. Jensen, Ph.D.



*Vitria Adisetiyo, Ph.D.*

"MRI relaxation rates are the more conventional way to measure brain iron, but they are not very specific," Dr. Adisetiyo said. "We added MFC because it offers more refined specificity."

The results showed that the 12 ADHD patients who had never been on medication had significantly lower MFC than the 10 ADHD patients who had been on psychostimulant medication or the 27 typically developing children and adolescents in the control group. In contrast, no significant group differences were detected using relaxation rates or serum measures. The lower brain iron levels in the non-medicated group appeared to normalize with psychostimulant medication.

MFC imaging's ability to noninvasively detect the low iron levels may help improve ADHD diagnosis and guide optimal treatment. Noninvasive methods are particularly important in a pediatric population, Dr. Adisetiyo noted.

"This method enables us to exploit inherent biomarkers in the body and indirectly measure dopamine levels without needing any contrast agent," she said.

If the results can be replicated in larger studies, then MFC might have a future role in determining which patients would benefit from psychostimulants—an important consideration because the drugs can become addictive in some patients and lead to abuse of other psychostimulant drugs like cocaine.

"It would be beneficial, when the psychiatrist is less confident of a diagnosis, if you could put a patient in a scanner for 15 minutes and confirm that brain iron is low," she said. "And we could possibly identify kids with normal iron levels who could potentially become addicts."

Along with replicating the results in a larger population of patients, the researchers hope to expand their studies to look at the relationship between cocaine addiction and brain iron.

Other co-authors are F. Xavier Castellanos, M.D., Adriana Di Martino, M.D., Kevin M. Gray, M.D., Els Fieremans, Ph.D., Ali Tabesh, Ph.D., and Rachael L. Deardorff, M.S.

###

Note: Copies of RSNA 2013 news releases and electronic images will be available online at [RSNA.org/press13](http://RSNA.org/press13) beginning Monday, Dec. 2.

RSNA is an association of more than 53,000 radiologists, radiation oncologists, medical physicists and related scientists, promoting excellence in patient care and health care delivery through education, research and technologic innovation. The Society is based in Oak Brook, Ill. ([RSNA.org](http://RSNA.org))

Editor's note: The data in these releases may differ from those in the published abstract and those actually presented at the meeting, as researchers continue to update their data right up until the meeting. To ensure you are using the most up-to-date information, please call the RSNA Newsroom at 1-312-949-3233.

For patient-friendly information on MRI, visit [RadiologyInfo.org](http://RadiologyInfo.org).

