

---

## RSNA Press Release

### Asymptomatic Atherosclerosis Linked to Cognitive Impairment

Released: November 25, 2014

Chicago — In a study of nearly 2,000 adults, researchers found that a buildup of plaque in the body's major arteries was associated with mild cognitive impairment. Results of the study conducted at the University of Texas (UT) Southwestern Medical Center will be presented next week at the annual meeting of the Radiological Society of North America (RSNA).

"It is well established that plaque buildup in the arteries is a predictor of heart disease, but the relationship between atherosclerosis and brain health is less clear," said

Christopher D. Maroules, M.D.,

radiology resident at UT Southwestern Medical Center in Dallas. "Our findings suggest that atherosclerosis not only affects the heart but also brain health."



*Christopher D. Maroules, M.D.*

#### At A Glance

- A large-scale study found a relationship between atherosclerosis and deterioration of brain health.
- Atherosclerosis is a condition in which fat, cholesterol and other substances collect in the arteries.
- Individuals with higher levels of internal carotid wall thickness were up to 21 percent more likely to have cognitive impairment.

Atherosclerosis is a condition in which fat, cholesterol and other substances collect in the arteries, forming a substance called plaque that can build up, limiting blood flow. It can occur in any artery of the body, including the carotid, which supplies blood to the brain, coronary arteries and the aorta, which carries oxygenated blood from the heart through the abdomen to the rest of body.

In the study, researchers analyzed the test results of 1,903 participants (mean age, 44 years) in the Dallas Heart Study, a multiethnic population-based study of adults from Dallas County, Texas. The participants included both men and women who had no symptoms of cardiovascular disease.

Study participants completed the Montreal Cognitive Assessment (MoCA), a 30-point standardized test for detecting mild cognitive impairment, and underwent magnetic

resonance imaging (MRI) of the brain to identify white matter hyperintensity (WMH) volume. Bright white spots known as high signal intensity areas on a brain MR images indicate abnormal changes within the white matter.

"Increased white matter hyperintensity volume is part of the normal aging process," Dr. Maroules explained. "But excessive WMH volume is a marker for cognitive impairment."

Study participants also underwent imaging exams to measure the buildup of plaque in the arteries in three distinct vascular areas of the body: MRI to measure wall thickness in the carotid arteries and the abdominal aorta, and computed tomography (CT) to measure coronary artery calcium, or the amount of calcified plaque in the arteries of the heart.

Using the results, researchers performed a statistical regression to correlate the incidence of atherosclerosis and mild cognitive impairment. After adjusting for traditional risk factors for atherosclerosis—including age, ethnicity, male sex, diabetes, hypertension, smoking and body mass index—they found independent relationships between atherosclerosis in all three vascular areas of the body and cognitive health, as measured by MoCA scores and white matter hyperintensity volume on MR images.

Individuals in the highest quartile of internal carotid wall thickness were 21 percent more likely to have cognitive impairment as measured by a low MoCA score. An increasing coronary artery calcium score was predictive of large white matter intensity volume on MRI.

"These results underscore the importance of identifying atherosclerosis in its early stages, not just to help preserve heart function, but also to preserve cognition and brain health," Dr. Maroules said.

Dr. Maroules said the MRI and CT imaging techniques provide valuable prognostic information about an individual's downstream health risks.

"Plaque buildup in blood vessels throughout the body offers us a window into brain health," he said. "Imaging with CT and MRI has an important role in identifying patients who are at a higher risk for cognitive impairment."

Co-authors on the study are Jarett Berry, M.D., Amit Khera, M.D., Heidi Rossetti, Ph.D., Kirstine Carter, Ph.D., Mohit K. Gupta, M.D., Ronald M. Peshock, M.D., Roderick McColl, Ph.D., Colby Ayers, M.S., Nagina Malguria, M.B.B.S., Suhny Abbara, M.D., and Kevin S. King, M.D.

###

Note: Copies of RSNA 2014 news releases and electronic images will be available online at [RSNA.org/press14](https://www.rsna.org/press14) beginning Monday, Dec. 1.

RSNA is an association of more than 54,000 radiologists, radiation oncologists, medical physicists and related scientists, promoting excellence in patient care and health care delivery through education, research and technologic innovation. The Society is based in Oak Brook, Ill. ([RSNA.org](https://www.rsna.org))

Editor's note: The data in these releases may differ from those in the published abstract and those actually presented at the meeting, as researchers continue to update their data right up until the meeting. To ensure you are using the most up-to-date information, please call the RSNA Newsroom at 1-312-791-6610.

For patient-friendly information on MRI and CT, visit [RadiologyInfo.org](https://www.RadiologyInfo.org).